

SUGGESTED APPLICATION AND PROTOCOL

**** Your Protocol and Application practice may vary based on your plant condition ****

We strongly suggest a **cocktail application** for best results. Standalone application is on a case by case basis and consulting with an Organnabis Technical Advisor (OTA) is recommended.



PREVENTATIVE MAINTENANCE

Dilution Rate: 1 ½ oz. Buddapest + ½ oz. pHlourish + 1 Gal. Water (Temperature not below 65 deg F)

Frequency: Day 1: Cocktail and apply Buddapest with pHlourish. Day 4: Apply pHlourish alone.

Application Method: Apply using atomizer, misting system or other spraying apparatus. Make sure that leaves are fully wet including the underside.

INDOOR: Apply at end of light cycle or by simply turning lights off. Wait until leaves are dry before turning lights on.

OUTDOOR: Dusk or Dawn (preferred). Avoid applying under direct light (This includes sunshine).

INFESTATIONS

Dilution Rate: 2 oz. Buddapest + ¼ oz. pHlourish + 1 Gal. Water (Temperature not below 65 deg F)

Occasional Severe: 3.5 oz. Buddapest + ¾ oz. pHlourish + 1 Gal. Water

Note: Higher concentration rates may be called for to treat severe infestation. However, no higher than 4oz./gal for Buddapest and no higher than 1oz/gal for pHlourish to avoid possible phytotoxicity.

Frequency: Apply 3 consecutive days, skip day 4, then once again on day 5.

Application Method: Apply using atomizer, misting system or other spraying apparatus. Make sure that leaves are fully wet including the underside.

INDOOR: Apply at end of light cycle or by simply turning lights off. Wait until leaves are dry before turning lights on.

OUTDOOR: Dusk or Dawn (preferred). Avoid applying under direct light (This includes sunshine).

**** IMPORTANT ****

Shake the bottles of Buddapest and pHlourish very well before diluting with water and again before each application. Water temp not below 65 deg F.

Organnabis Technical Advisor Hotline: (720) 372-0513

Email: info@organnabis.us

www.organnabis.us



ORGANNABIS.US